

Children's After School Center



MEDICAL DECLARATION STATEMENT FOR SCHOOL-AGE CHILD CARE

(AND/OR FOR CHILDREN ENROLLED IN PUBLIC OR PRIVATE SCHOOL)

CHILD'S NAME:	DATE OF BIRTH:	GRADE IN SEPTEMBER:

HEALTH STATEMENT (CHECK ONE)

- My child is in good health and can participate in the normal activities of the program and has no conditions or special needs that require special accommodations.
- My child can participate in the normal activities of the program but has conditions or special needs that require special accommodations as indicated below.

SCHOOL-AGE CHILD'S SPECIAL CONDITIONS OR NEEDS REQUIRING SPECIAL ACCOMMODATIONS

Please list any allergies, medical conditions, including chronic health problems (such as asthma, seizures), behavioral disorders, special needs, etc.

PARENT/GUARDIAN SIGNATURE:	DATE:

PARENTAL AUTHORIZATION FOR EMERGENCY TREATMENT

Name Of Child:	Birthdate:	Enrollment Date:
----------------	------------	------------------

PARENT/GUARDIAN INFORMATION	<input type="checkbox"/> PARENT/GUARDIAN # 1		<input type="checkbox"/> PARENT/GUARDIAN # 2	
	Name:		Name:	
	Relationship:		Relationship:	
	Cell Phone:		Cell Phone:	
	Home Phone:		Home Phone:	
	Home Address:		Home Address :	
	Employer Name:		Employer Name:	
	Employer Phone:		Employer Phone:	
	E-Mail Address:		E-Mail Address:	

EMERGENCY CONTACTS	Persons authorized to pick up your child and/or contact in case of emergency if neither parent is available to assume responsibility for the child.					
	Contact Name #1:		Contact Name #2:		Contact Name #3:	
	Relationship:		Relationship:		Relationship:	
	Cell Phone:		Cell Phone:		Cell Phone:	
	Home Phone:		Home Phone:		Home Phone:	
	Employer Phone:		Employer Phone:		Employer Phone:	

CUSTODY	Name of person PROHIBITED from picking up your child:	
	If a non-custodial parent has been denied access, or granted limited access, to the child by a court order, please submit documentation to this effect for the center to maintain a copy on file, and to comply with the terms of the court order.	

MEDICAL INFORMATION	Child's Health Care Provider:	
	Health Care Provider Phone:	
	Health Care Provider Address:	
	Name Of Insurance Company/Hmo:	
	Group #:	
	Identification #:	
	Subscriber's Name On Insurance Card:	
	Known Allergies (including medication):	
	Medication My Child Is Taking:	
	List Special Conditions, Disabilities, Medical/Physical Restrictions, Medical Information For Emergency Situations:	

AUTHORIZATION FOR EMERGENCY MEDICAL TREATMENT	
As the parent(s)/ legal guardian(s) of the above named child, I (we) attest that the information above is correct. I (we) authorize the child care center staff to obtain emergency treatment for my child and understand that I (we) shall be promptly notified.	

Parent/Guardian Signature #1:	Date:	Parent/Guardian Signature #2:	Date:
-------------------------------	-------	-------------------------------	-------



PLACE
PICTURE
HERE

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: [] Yes (higher risk for a severe reaction) [] No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

[] If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for ANY symptoms.

[] If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING: SEVERE SYMPTOMS



LUNG

Short of breath, wheezing, repetitive cough



HEART

Pale, blue, faint, weak pulse, dizzy



THROAT

Tight, hoarse, trouble breathing/swallowing



MOUTH

Significant swelling of the tongue and/or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION of symptoms from different body areas.



- 1. INJECT EPINEPHRINE IMMEDIATELY.**
- 2. Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy/runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea/discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: [] 0.15 mg IM [] 0.3 mg IM


Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

EPIPEN® AUTO-INJECTOR DIRECTIONS

1. Remove the EpiPen Auto-Injector from the clear carrier tube.
2. Remove the blue safety release by pulling straight up without bending or twisting it.
3. Swing and firmly push orange tip against mid-outer thigh until it 'clicks'.
4. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove auto-injector from the thigh and massage the injection area for 10 seconds.



ADRENALICK® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR DIRECTIONS

1. Remove the outer case.
2. Remove grey caps labeled "1" and "2".
3. Place red rounded tip against mid-outer thigh.
4. Press down hard until needle enters thigh.
5. Hold in place for 10 seconds. Remove from thigh.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____

PHONE: _____

NAME/RELATIONSHIP: _____

PHONE: _____